

Magnetic Surface Visualizations in the Columbia Non-neutral Torus

Paul W. Brenner, Thomas Sunn Pedersen, John W. Berkery, Quinn R. Marksteiner, Michael S. Hahn

Abstract—Visualizations of magnetic surfaces are a valuable diagnostic in the Columbia Non-neutral Torus (CNT). CNT is a compact stellarator, which is currently being used to study non-neutral plasmas confined on magnetic surfaces. The full three-dimensional shapes of magnetic surfaces created by CNT's simple four circular coil geometry are readily visualized using an electron beam and neutral gas. These visualizations are useful for probe alignment, confirmation of the magnetic surface topology, and were necessary for the recent installation of a conducting boundary conforming to the last closed magnetic surface.

Index Terms—non-neutral, stellarator, magnetic surfaces, visualization

AS a stellarator, the Columbia Non-neutral Torus (CNT) offers unique benefits for the study of non-neutral plasmas [1]. Plasmas of various degrees of neutrality can be studied because no external electric field is required for confinement. Steady state low density plasmas can be confined without internal currents. Long confinement times are expected as a result of the the large space charge electric field [2].

Magnetic surfaces in CNT are created using a simple four circular coil design with modest tolerances. Thus design, optimization, construction, and installation were relatively fast and inexpensive [1]. Currently the confinement [2], stability [3], and equilibrium [4] of pure electron plasmas are being studied in CNT.

Pure electron plasmas in CNT are usually studied with pressures in the 10^{-9} Torr range to minimize the effect of electron neutral collisions. However, by backfilling to a pressure above 10^{-5} Torr and emitting a beam of electrons from an electron gun it is possible to visualize the complete three-dimensional shape of the magnetic surfaces (Fig.1). The electron gun is a heated-biased filament surrounded by a grounded metal cap with a hole in the end. Emitted electrons are accelerated out of the cap and travel along the magnetic surface with enough energy to excite and ionize the neutral background gas which then emits visible light. Brightness increases with background pressure at the cost of reduced electron beam path length.

Although magnetic surface topology was originally confirmed [1] using the standard electron beam-fluorescent rod technique [5], the visualization method was also used to demonstrate CNT's magnetic surfaces. An example of such




Fig. 1. A photograph of a surface visualization made using an electron gun taken with 12 second exposure on a digital camera.

a visualization at 5×10^{-5} Torr with a 200 eV electron beam is shown in Fig.1. These glowing magnetic surfaces are bright enough to be seen by the naked eye.

Recently, a conducting boundary conforming to the last closed flux surface (LCFS) was installed. The conducting boundary improves confinement by setting the electrostatic potential at the LCFS, reducing the mismatch between equipotential and magnetic surfaces in the confining volume [6].

The conducting boundary has also improved visualizations. The e-gun's grounded metal cap collected the majority of emitted electrons. The cap is no longer required because the grounded conducting boundary is sufficient to accelerate electrons off the bare filament. Therefore, more emitted electrons are free to travel along the surface. This results in significantly brighter visualizations that proved necessary for proper installation of the mesh.

Proper alignment of the conducting boundary is necessary to ensure the segments are not limiting the plasma. Without visualizations, alignment would be a complicated process. However, visualizations make misalignments quite obvious (Fig. 2(a)). Since installation, adjustments have been made with this alignment method so that the conducting boundary encloses a large confining volume (Figs. 2(b) and 2(c)).

Manuscript received January 10, 2008; revised ???????? 1, 2008. Supported by the Fusion Energy Sciences Postdoctoral Research Program of the U.S. Department of Energy and by the NSF CAREER program, Grant No. NSF-PHY-04-49813

The authors are with the Department of Applied Physics and Applied Math, Columbia University, 500 W. 120th st, New York, NY, 10027.(e-mail:pwb2103@columbia.edu)

Digital Object Identifier S XXXX-XXXXXXX-X




Fig. 2. Visualization photographs taken with 4 second exposure on a digital camera. (a) A glowing field line is plainly visible intersecting the conducting boundary prior to alignment. (b)(c) Large outer surfaces glowing inside the conducting boundary without intersecting it.

In conclusion, these visualizations are uniquely useful for demonstration of magnetic surfaces, probe alignment, and were critical for alignment of a recently installed boundary.

REFERENCES

- [1] T. S. Pedersen, J. Kremer, R. Lefrancois, Q. Marksteiner, N. Pomphrey, W. Reiersen, F. Dahlgreen, and X. Sarasola, "Construction and initial operation of the Columbia Nonneutral Torus," *Fusion Science and Technology*, vol. 50, pp. 372–381, 2006.
- [2] J. Berkery, T. S. Pedersen, J. Kremer, Q. Marksteiner, R. Lefrancois, M. Hahn, and P. W. Brenner, "Confinement of pure electron plasmas in the Columbia Non-neutral Torus," *Physics of Plasmas*, vol. 14, no. 6, 2007.
- [3] Q. Marksteiner, T. S. Pedersen, J. Berkery, M. Hahn, J. Mendez, and H. Himura, "Observations of an ion driven instability in non-neutral plasmas confined on magnetic surfaces," *accepted to Physical Review Letters*, 2007.
- [4] J. Kremer, T. S. Pedersen, Q. Marksteiner, and R. Lefrancois, "Experimental confirmation of stable, small Debye length pure electron plasma equilibria in a stellarator," *Physical Review Letters*, vol. 97, 2006.
- [5] R. Jaenicke, E. Ascasibar, P. Grigull, I. Lakicevic, A. Weller, M. Zippe, H. Hailer, and K. Schworer, "Detailed investigation of the vacuum magnetic surfaces on the W7-AS stellarator," *Nuclear Fusion*, vol. 33, no. 5, pp. 687–704, 1993.
- [6] R. Lefrancois, T. S. Pedersen, A. Boozer, and J. Kremer, "Numerical investigation of three-dimensional single-species plasma equilibria on magnetic surfaces," *Physics of Plasmas*, vol. 12, p. 072105, 2005.